

# Alka Seltzer Poppers: An Interactive Exploration

A. M. Sarquis\* and L. M. Woodward\*

Center for Chemical Education, Miami University Middletown, Middletown, OH 45042-3497

Few teachers will argue that it is difficult to interactively engage students in the experimental nature of chemistry through traditional lecture and exam methods. The following student challenge is offered as an alternative to traditional methods of assessing student learning.

We have found that using questioning strategies to involve students in actually doing and analyzing experimental design can offer useful insights to instructors and students alike. Assessment is based not only on getting a right answer but also on the logic of the students' approaches, the methods and documentation presented by the students, and their understanding of the system, based in part on their answers to the questions.

The following interactive exploration has been used when discussing topics such as chemical reactions, gases, solubility, and kinetics.

## Student Challenge

Alka Seltzer is sometimes taken for acid indigestion. This over-the-counter medication is taken differently from most others in that it is added to water before ingesting, rather than using water to "wash it down". Let's investigate this addition of Alka Seltzer to water. What are the effects of changing the volume and the temperature of the water used in the investigation?

### Materials

- 5 (or more) Alka Seltzer tablets
- clear container such as a plastic cup or glass
- room-temperature water
- (optional) 1/4-cup measure
- one 35-mm film canister with lid
- teaspoon measure
- tablespoon measure
- ice-cold water
- very hot tap water

### Safety

**CAUTION:** Goggles are recommended to protect your eyes from flying debris that may occur in this activity. Do not use water hot enough to scald.

### Exploration

**Step 1:** Determine and record the active ingredients of Alka Seltzer tablets.

**Step 2:** Add about 1/4 cup of room-temperature water

into a clear container. Add an Alka Seltzer tablet to the water and observe. Keep the solution for use in Step 3. Describe what happens. What role does water play in this system? What is the identity of the most obvious product of this chemical change? Write the chemical equation that describes the reaction that has occurred.

**Step 3:** Place 1 tablespoon of the solution from Step 2 into the film canister. Add half an Alka Seltzer tablet to the canister and immediately put on the lid securely. Hold onto the sides of the container, so that the lid is pointed upward and away from your face. What happens? Explain your observations.

**Step 4:** Design and perform an investigation to determine if the observations in Step 3 are reproducible. What did you do? What did you find out?

**Step 5:** Repeat Step 3 with room-temperature water that has not had Alka Seltzer dissolved in it already. How do the results compare with those in Step 3?

**Step 6:** Rinse out the canister and place 1 teaspoon of room-temperature water in it. Add half an Alka Seltzer tablet to it and immediately put on the lid as before. Compare your observations with those in Step 5 and explain any similarities and differences.

**Step 7:** What do you think would happen if you added 2 tablespoons of water to the canister before adding the Alka Seltzer? Try this variation and explain your observations.


**Step 8:** Design an experiment to determine the impact of changes in the temperature of the water on the system. Do not use water hotter than hot tap water or colder than ice water. Describe what you did and your observations. Explain what you observed.

**Step 9:** Which trial is the control, the one to which the other trials are compared? Why is the number of variables that are changed in any one trial important? What other variables might be studied in this system?

## Expected Student Responses


**Step 1:** The active ingredients in Alka Seltzer are sodium bicarbonate, citric acid, and aspirin (acetylsalicylic acid).

**Step 2:** The solution bubbles and the Alka Seltzer tablet disappears. The water dissolves the reactants and facilitates the reaction. The most obvious product is the gas, which is carbon dioxide. The reactions are acid-base neutralization reactions between the base, sodium bicarbonate, and each of the two acids. With citric acid the reaction is:



With aspirin (acetylsalicylic acid), the reaction is:

\*Email: sarquiam@muohio.edu and woodwal@muohio.edu.



**Step 3:** As the carbon dioxide is produced, pressure builds up inside the canister until, after a few seconds, it is great enough to force the lid off.

**Step 4:** Students should repeat the experiment several more times to check for reproducibility. They should measure the time it took for the lid to be forced off for comparison. They should find that the results are reproducible to within a second or two.

**Step 5:** In Step 3, the water was already saturated with carbon dioxide so all the gas produced went toward increasing the pressure within the canister. In Step 5, some of the carbon dioxide gas produced dissolved in the water so it took longer to generate enough gas to cause the top to pop.

**Step 6:** With one teaspoon of water instead of one tablespoon, it takes longer for the top to pop off the canister. There are several factors to consider. There is less water so the tablet may dissolve less rapidly and less of the carbon dioxide dissolves in the water. There is also more air space so it takes longer to produce enough gas to increase the pressure enough to pop the top.

**Step 7:** Two tablespoons of water nearly fill the canister. There is more water for the tablet and the carbon dioxide to dissolve in. There is very little space for gases, so a relatively small amount of gas will get the pressure high enough to force the top off.

**Step 8:** Step 5 should be repeated with different temperatures of water and comparisons made. The reaction rate increases (and gas solubility decreases) with increasing temperature so the top pops fastest with the hottest water and slowest with the coldest water.

**Step 9:** The trial in Step 5 functions as the control. One variable was altered in each trial (although this sometimes had several opposing effects). If more than one variable is altered, it is difficult or impossible to determine which one accounted for observed changes. Another variable that could be tested is the amount of Alka Seltzer tablet used.

### Answer to Challenge

Water acts as a solvent to dissolve the active ingredients (sodium bicarbonate and the two acids) so that they can react. The rate of the pop is directly proportional to both the temperature of the system and to the volume of the water used.

### Acknowledgments

We wish to acknowledge the contribution of Jill Selzer, who introduced us to the popping canister phenomenon through her term project in our Teaching Science with TOYS course, and the National Science Foundation Division of Undergraduate Education for its support of the curriculum development effort (grant #9055448).